



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

developed and the most posterior ones minute, so the personal equation enters into their count somewhat. Aside from this there is sufficient variation to bridge the difference, 42 to 51 counted in 8 specimens of *crysos* from North Carolina and Florida, in the American Museum of Natural History.

The longer pectoral credited to *caballus* and *pisquetus* is an age character. As is common in carangin species the pectoral increases irregularly with age and becomes more falcate. *Caranx crysos* over 125 mm. in length to base of caudal are sufficiently like the adult to be recognized for what they are at a glance. One of 129 mm., Havana Market, has pectoral 3.6 in length, .97 in head; 5 of 150 to 165 mm. from North Carolina and Florida, 3.0 to 3.2 (average 3.12) in length, .86 to .94 (average .91) in head; 3 of 215 to 235 mm. from North Carolina, 3.0 to 3.2 (average 3.07) in length, .84 to .86 (average .85) in head; one of 311 mm. from Brazil, 2.7 in length, .76 in head; one of 430 mm. from Cape San Lucas, 3.1 in length, .84 in head.

Caranx crysos (Mitchill), New York, was based on small specimens; large ones appear to be rare to the north.

J. T. NICHOLS,
New York, N. Y.

A BLACK PITUOPHIS

My friend, Dr. H. P. Loeding of Mobile, Ala., recently sent me for examination a large Pituophis, remarkable for being uniformly black above and below, except for a little rusty color on the anterior part of the head and flecks of rusty on the ends of some of the ventral scales.

This is apparently the first example of its genus to be reported from Alabama, and, so far as I know, there are no records for Georgia, Mississippi and Louisiana. This specimen was found dead on the

Hall's Mill Road, in the vicinity of high, sandy hills near Hall's Mill Creek, about 14 miles southwest of Mobile. A second specimen, which Dr. Loeding informs me is like this one, was taken alive at Grand Bay, 26 miles southwest of Mobile, by Mr. E. D. King, Jr. The latter, a female, was kept in confinement for over a week, but refused to eat, so was preserved and deposited in the Charles Mohr Museum in Mobile.

As Dr. Loeding and his friends have been collecting reptiles in the vicinity of Mobile for many years, it would seem as if the ordinary patterned form of *Pituophis* would have been found if it occurred at all commonly in Mobile County. On the other hand the black form was not found until this past season. Possibly, however, it has been confused in the field with some of the more common large black snakes, as *Drymarchon corais couperi* and *Coluber constrictor constrictor*.

The finding of two black specimens of *Pituophis* so far apart as twelve miles, and the absence of records for normally colored individuals, suggest the exclusive occurrence in this region of a black phase of the Bull Snake.

That all the North American bull snakes are very closely related can hardly be doubted. In fact it appears that only color pattern can be relied upon to distinguish them with certainty. We would expect, therefore, to find each species of *Pituophis* directly related to the one inhabiting the adjacent range. We would not then look for a black form occupying a range between the ranges of two closely allied patterned forms. In the United States National Museum there are two examples of *Pituophis* (No. 10,363) from Murphy (near Knoxville), Tenn., patterned like the eastern species, *P. melanoleucus*. Our black phase could therefore not extend farther north than this point, and it is very likely that it will prove to be restricted to the extreme South,

where it is now found. If the Gulf Coast may be regarded as in general unfavorable to the bull snakes, we may readily understand how a local color phase may have become established in a limited region of favorable habitat.

As there are no other specimens of *Pituophis* on record from this southern tier of states, and since it is chiefly on color characters that the nearest species, *P. sayi* and *P. melanoleucus* are distinguished, it is rather difficult to assign this black form definitely to either one. The two specimens from Murphy, Tenn., are undoubtedly *P. melanoleucus*. This is perhaps the most western definite record for the eastern species. How much farther west it may extend can only be conjectured. Eastern records for *P. sayi* seem to be limited to Illinois, but numerous western species of reptiles are known east of the Mississippi River only in Illinois and Indiana. These facts, and the apparently unfavorable habitat of the lower Mississippi River region, suggest that *P. melanoleucus* may occupy all the favorable areas in the southern states as far west as the Mississippi River. In that case our black specimens may be regarded as a local phase of *P. melanoleucus*.

Carination and scutellation are admittedly unsatisfactory in distinguishing *P. sayi* from *P. melanoleucus*. Our black example has five rows of smooth scales on the sides, and Dr. Loeding informs me that the other specimen has the same number. This, while within the extremes of *P. melanoleucus*, comes perhaps closer to the average for *P. sayi*. Similarly, the rostral dividing the internasals for only two-thirds of their width rather suggests *P. sayi*. However, the Tennessee specimens mentioned above, are practically like this black individual in carination and shape and size of rostral, while definitely *P. melanoleucus* in pattern. It therefore seems more satisfactory to assign these black specimens provisionally to *P. melanoleucus*.

The finding of more examples of *Pituophis* from these southern states will be awaited with much interest.

Following is a description of the black example from 14 miles southwest of Mobile, now deposited, through the kindness of Dr. Loeding, in the United States National Museum (No. 62,340):—

Ventrals, 225; anal, single and entire; 57 divided caudals; upper labials, 8 on each side, lower labials, 13 on the left side and 15 on the right; one preocular on each side; 4 postoculars on the left side and probably 4 on the right; about 4 temporals in the first row; rostral dividing the internasals for two-thirds of their width; maximum number of scale rows, 29, anteriorly, 27; posteriorly, 21; keels on dorsal scales prominent above, progressively fainter on the sides, descending as low as the sixth row anteriorly and the third row posteriorly. Total length, 1,800 millimeters; tail length, 221 millimeters. Sex, female.

The coloration (by reference to Ridgway's Color Standards and Nomenclature) is as follows: Above, fuscus black; below, slate color; on the head, between the parietals and the rostral, and including the upper labials, most of the scales having in their centers a development of orange-cinnamon mixed with the fuscus black; occasional ventral scales, except on the anterior portion of the body, with flecks of perhaps an ivory yellow, or lighter, near their ends; and along the sides of the tail and near its end, on most of the subcaudal scutes, some lighter coloration showing faintly through the black.

FRANK N. BLANCHARD,
University of Michigan.

PILOT SNAKE SWALLOWING NEST EGG

I was greatly interested in the account recently published in Copeia of a Pilot Snake, *Elaphe obso-*